

**The ameliorative effect of *alkanna orientalis* extract against Horned viper venom renal
lipogram and cholinergic toxicity**

Mahmoud. M. Nasser¹; Abd El Aziz. A. Diab²; Ahmed. L. S. Hashem¹; Samih. I. El
DAHmy³ and Eman. M. E. Mohallal¹

1 physiology department, Desert research centre, Cairo, Egypt.11753

2 Zoology department, faculty of science, Zagazig university, Egypt.

3 pharmacognocny department, faculty of pharmacy, Zagazig university, Egypt.

Corresponding author: Nasser_mahmoud55@yahoo.com

ABSTRACT : The present study designed to evaluate the toxicity of Horned viper/*Cerastes cerastes* (*Cc*) venom on male albino rats and the possible ameliorative role of *Alkanna orientalis* (*Ao*) extract against venom toxicity. Forty-eight adult male albino rats (180 ± 200 g) divided into 6 groups. Group I injected intraperitoneal with physiological saline (100µl). Group II injected intraperitoneal with *Ao* extract (250 mg/kg/BW). Group III injected with 1/10 LD₅₀ of *Cc* venom (0.435 mg/kg/BW). Group IV injected intraperitoneal with *Cc* venom (0.435 mg/kg/BW) then immediately injected with *Ao* extract (250 mg/kg/BW). Group V was given *Ao* extract orally (250 mg/kg/b.wt) then after 2 hours injected with *Cc* venom (0.435 mg/kg/BW) Group VI was injected with *Cc* venom (0.435 mg/kg/BW) then was injected immediately with immunoglobulin (300µl). Group III showed a significant increase in serum creatinin, urea, uric acid as well as a significant decrease in serum total cholesterol, triglyceride, HDL-cholesterol, LDL-cholesterol, glucose level, calcium and cholinesterase when compared with control group. As well as, *Ao* extract when given intraperitoneal immediately or orally 2 hours before *Cc* venom as antidote, it minimizes the changes of previous biochemical parameters. groups IV and V when compared with group VI, it showed that *Ao* extract considered more effective antidote than antivenom immunoglobulin for *Cc* venom renal, lipogram and cholinergic toxicity. In conclusion, *Ao* extract showed a new therapeutic and prophylaxis agents against *Cc* venom renal, lipogram and cholinergic toxicity.

KEYWORDS: *Cerastes cerastes*; *Alkanna orientalis*; family *Boraginaceae*; venom toxicity; haematological effect.

Date of Submission: 01-08-2022

Date of acceptance: 23-09-2022

I. INTRODUCTION

Snake's envenomation is a critical public medical health problem [1] resulting in high mortality and morbidity rate [2]. Venomous snakes documented as the most dangerous poisoning animals all over the world and the fatality of its bites depending on the injected venom amount, location of the bite, the size of the victim, the snake species and the time between the bite and the injection of anti-venom [3]. The toxic effects of the venoms result from both protein and non-protein components which have distinct actions [4].

Cerastes cerastes venom was reported to cause local tissue damage, that lead to the extra cellular matrix disruption as well as, the basement membrane disruption that covering the body vessels and blood capillaries [5].

[6] Reported that, the cell membrane permeability changes due to the venom toxicity that resulting in the internal cell content discharge, oxidative stress and it cause an inflammatory response. Therefore, anti-inflammatory and antioxidant drugs may be important in the treatment of snake bite. In addition, the neurological toxic effect of snakebites such as neuromuscular paralysis was documented by [7].

The usage of anti-venom can neutralize the venom as well as, it can prevent further damage but it cannot restore the damage that was already occurred [8]. Therefore, it is seriously to find out new natural or artificial venom inhibitors, which can compete the anti-venom therapy or complement the anti-venom action especially, in the local tissue damage restoring [9]. Glycosaminoglycans, polysaccharides, flavonoids and alkaloids was documented as venom spreading factors inhibitors [10]. Herbs and plants were recorded as pharmacological ingredients source that represent about 25% of the recently used drugs [11]. Because of the low-cost traditional medicinal plants, it is commercially used for diseases treatment especially in low-income population's countries [12].

Boraginaceae family includes several plants that widely distributed in mild and tropical regions and it was classified into 200 genus and 2000 species, most of these family members have important medicinal usage. Some plants of *Boraginaceae* family documented as snakebite antidote such as, *Argusia argentea* extract is effective in *Trimeresurus flavoviridis* bites [13]. Whole the plant of *Carmona retusa* reported as antidote paste for *Daboies russelii* bites [14]. Rosmarinic acid from *cordial verbenacea* extract is effective against *Bothrops jararacussu* bites [15].

Alkanna orientalis and *Echium humil*, belong to *Boraginaceae* family rich with stearidonic acid (SDA), alpha-linolenic acid (ALA) and gamma-linolenic acid (GLA) in addition to rosmarinic acid (RA). All previous ingredient was reported to be a powerful anti-inflammatory and antioxidant effect so; they can restore the *Cerastes cerastes* venom toxicity [16] and [17].

Rosmarinic was reported as active ingredient of *Alkanna orientalis* that showed antioxidant, anti-inflammatory and antibacterial effect [18]. As well as, **rosmarinic** acid able to inhibit the snake venom phospholipases A2 and metalloproteinase from *Cordia verbenacea* and *B. jararaca* [19].

On the same pattern, [20] reported flavonols fractions from aerial parts of *Alkanna orientalis*. Flavanoids are one of the foremost plant components that work against PLA2 and lipooxygenase, they showed anti-allergic, anti-inflammatory and enzyme inhibiting properties. Flavanoid weakly inhibits the group I PLA2 from *Naja naja* and strongly inhibits the group II PLA2 from *Vipera* [21].

II. AND METHODS II.MATERIALS

Experimental animals

Eighty eight male albino rats, weight range 180–200 gm, obtained from faculty of science animal house, Zagazig University. These rats quarantined 15 days before the experiments carried, settled in plastic cages, fed a standard diet and water, exposed to a 12 h light/dark cycle, and maintained at temperature range 22±2°C. All Institutional and National Guidelines for the care and use of animals were followed.

Cerastes cerastes venom

25 *Cerastes cerastes* vipers obtained from El Kharga (New Valley), settled in cages at in desert research center physiology lab. Vipers were milked by using a rubber into a small beaker, and then the venom was rapidly frozen by using liquid nitrogen until the study start.

Cerastes cerastes venom LD₅₀

LD₅₀ of *Cerastes cerastes* venom was calculated according to the method that described by [22]. 40 adult male albino rats were divided into 5 groups (gp I-V), injected intraperitoneal (i.p.) with *Cerastes cerastes* venom with doses (2.0, 3.0, 4.0, 5.0, 6.0) mg/kg/BW for each group, respectively. 24 hours post injection the dead and alive rats in each group were recorded. The calculated LD50 was 4.35 mg / kg/BW.

Alkanna orientalis extraction

Alkanna orientalis extract was prepared according to [23] method for intraperitoneal administration. The plant obtained from Saint Katherine, South Sinai, Egypt, where it grow, and then it was left in air for drying away from direct sun heat. Then, we grinded the plant then immersed in 70% ethyl alcohol for 4 days. The extract filtered and concentrated at room temperature. The extract stored at 4°C until using in the study.

Antivenom

We purchased Anti-venom immunoglobulin from VACSERA CO. Agouza .Giza, Egypt.

Animal treatment schedule

Forty eight male albino rats were divided into six groups (n=8/group), as the following:

Group I (control group):

Rats were injected (i.p.) with 100µl of 0.09% physiological saline

Group II (*Alkanna orientalis* treated group):

Rats were injected (i.p.) with *Alkanna orientalis* extract (250 mg/kg/BW) dissolved in 100µl distilled water.

Group III (*Cerastes cerastes* venom treated group):

Rats injected (i.p.) with *Cerastes cerastes* venom (0.435 mg/kg BW) that represent 1/10 LD₅₀, dissolved in 100µl of 0.09% physiological saline.

Group IV (*Cerastes cerastes* venom + *Alkanna orientalis* extract) group:

rats injected (i.p.) with 1/10 LD₅₀ *Cerastes cerastes* venom (0.435mg/kg/BW) dissolved in 100µl of 0.09% physiological saline then, immediately injected (i.p.) with *Alkanna orientalis* extract (250 mg/kg/BW) dissolved in 100 µl distilled water.

Group V (a prophylaxis group):

Rats were given *Alkanna orientalis* extract orally (250 mg/kg/BW) then after 2 hours were injected (i.p.) with 1/10 LD₅₀ *Cerastes cerastes* venom (0.435 mg/kg BW) dissolved in 100µl of 0.09% physiological saline.

Group VI (*Cerastes cerastes* venom + antivenom):

Rats were injected (i.p.) with 1/10LD₅₀ of *Cerastes cerastes* venom (0.435 mg / kg/BW) dissolved in 100 µl of 0.09 % physiological saline, and then were immediately injected (i.p.) with 300 µl of antivenom immunoglobulin.

Blood sampling

In this study rats were sacrificed 2 hours after *Cerastes cerastes* venom injection and in group I & II, rats were sacrificed 2 hours after saline and *Alkanna orientalis* extract injection respectively, and blood samples were collected in sterile tubes without anticoagulant. The serum separated from clotted blood and used for the biochemical parameters determination.

Evaluation of some biochemical parameters

Biochemical parameters carried out by spectrophotometric analyzer (BT-260). Creatinin test measured according to method described by [24]. Serum urea, measured according to [25], Serum uric according to [26]. Serum calcium according to [27], Cholinesterase level according to [28], glucose according to method described by [29], Serum cholesterol According to method described by [30], serum triglycerides according to method described by [31] and Serum LDL and HDL-cholesterol was determined according to [32].

Statistical analysis

The obtained data were analyzed by the statistical analysis software (SAS–2013 program), for obtaining Mean values ±standard error. Subsequent multiple comparisons between the different groups were analyzed by Duncan's multiple comparison tests [33], values at ($P < 0.05$) were considered significant [34].

III. RESULTS

Kidney function parameters and cholinesterase

Table (1) illustrate the effect of *Cerastes cerastes* venom and the ameliorative role of *Alkanna orientalis* extract and antivenom on some kidney function parameters, calcium and cholinesterase in the different groups. Rats that injected with *Cerastes cerastes* venom (group III), showed a significant increase ($P < 0.001$) in level of serum creatinin, urea, uric acid and a significant decrease ($P < 0.001$) in serum calcium and cholinesterase level as compared with control group (group I). While, *Alkanna orientalis* extract (group II) did not show any significant effect in all previous parameters when compared with control group.

On the other hand, when *Alkanna orientalis* was injected (i.p.) immediately after *Cerastes cerastes* venom or given orally 2 hours before *Cerastes cerastes* venom as a prophylaxis dose, (groups IV and V, respectively), it was able to ameliorate the renal toxic effect of the venom as well as cholinergic toxicity. In addition, it showed a significant decrease ($P < 0.001$) in serum level of creatinin, urea, uric acid and a significant increase ($P < 0.001$) in serum calcium and cholinesterase as compared with venom treated group. On the same pattern, antivenom immunoglobulin showed a

significant ameliorative effect against the venom when given (i.p.) immediately after *Cerastes cerastes* crude venom injection.

Lipogram parameters and glucose level

Concerning to the effect of *Cerastes cerastes* venom and the ameliorative role of *Alkanna orientalis* extract and antivenom immunoglobulin on lipogram parameters and glucose level. Table (2) showed that, group III has a significant decrease ($P < 0.001$) in total cholesterol, triglyceride, HDL and LDL cholesterol as well as, serum glucose level in comparison with group I. While, *Alkanna orientalis* extract (group II) showed a significant decrease in triglyceride level and did not give any significant effect in the other parameters after 2 hours from (i.p.) injection of it as compared with control group as shown in Table (2). As well as, in groups IV and V *Alkanna orientalis* was able to ameliorate the toxic effect of the venom. It showed a significant increase ($P < 0.001$) in all previous parameters when compared with the venom treated group (group III). On the same pattern, antivenom immunoglobulin showed a significant ameliorative effect against the venom when given (i.p.) immediately after *Cerastes cerastes* crude venom injection. It showed a significant increase ($P < 0.001$) in all previous parameters when compared with the venom treated group (group III), as shown in Table (2).

These results showed that *Alkanna orientalis* extract when given as antidote for *Cerastes cerastes* venom renal, cholinergic and lipogram toxicity was effective hand by hand such as synthesized antibody (anti-venom immunoglobulin) and these results clearly appeared by comparing the findings of groups IV and V with group VI.

Table (1): effect each of *Cerastes cerastes* venom, *Alkanna orientalis* and their combination on some kidney functions, calcium and cholinesterase in male albino rats (mean \pm SE) (n=8).

groups	(mean \pm SE) of some kidney function parameters, calcim and cholinesterase (n = 8)				
	Creatinin mg\dl	Urea mg\dl	Uric acid mg\dl	CHE U\L	Ca mg\dl
1. Control group	0.78 \pm 0.09 ^b	43.75 \pm 2.2 ^d	4.26 \pm 0.23 ^d	3901 \pm 139 ^{ab}	10.48 \pm 0.4 ^a
2. Ao extract group	0.74 \pm 0.08 ^{bc}	45.75 \pm 3.6 ^d	4.53 \pm 0.22 ^{dc}	4004 \pm 111 ^a	10.45 \pm 0.2 ^a
3. Cc.venom group	1.41 \pm 0.05 ^a	85.5 \pm 4.1 ^a	8.80 \pm 0.15 ^a	2145 \pm 123 ^d	7.1 \pm 0.3 ^c
4. Cc.venom _{i.p} + Ao extract _{i.p} group	0.79 \pm 0.08 ^b	71.37 \pm 5.3 ^b	4.41 \pm 0.23 ^{cd}	3545 \pm 118 ^b	9.56 \pm 0.5 ^b
5. Ao extract _{oral} then 2h + Cc.venom _{i.p} group	0.71 \pm 0.07 ^{dc}	43.25 \pm 2.4 ^d	3.87 \pm 0.14 ^d	3568 \pm 133 ^b	10.56 \pm 0.6 ^a
6. Cc.venom _{i.p} +Antivenom _{i.p} group	0.79 \pm 0.04 ^b	63.62 \pm 2.3 ^c	5.73 \pm 0.29 ^b	2592 \pm 98 ^c	10.57 \pm 0.5 ^a

Means within the same column in each category carrying different litters are significant at ($P \leq 0.05$).

Where

Cc. Venom _____ *Cerastes cerastes venom*

Ao extract _____ *Alkanna orientalis extract*

I.p _____ *intraperitoneal*

Table (2): effect each of *Cerastes cerastes* venom, *Alkanna orientalis* and their combination on lipogram and glucose level in male albino rats (mean \pm SE) (n=8).

groups	(mean \pm SE) of lipogram and glucose parameters (n = 8)				
	cholesterol mg\dl	Triglyceride mg\dl	HDL cholestero mg\dl	LDL cholestero mg\dl	RBS mg\dl
1. Control group	99.0 \pm 2.5 ^a	80.1 \pm 5.2 ^a	30.2 \pm 1.1 ^b	56.3 \pm 2.1 ^c	75.8 \pm 2.3 ^a
2. <i>Ao</i> extract group	98.2 \pm 3.1 ^a	63.1 \pm 4.3 ^{cd}	30.1 \pm 0.9 ^b	55.3 \pm 1.6 ^c	76.6 \pm 3.1 ^a
3. <i>Cc.venom</i> group	80.5 \pm 4.2 ^c	66.5 \pm 3.2 ^c	23.1 \pm 1.3 ^c	50.0 \pm 1.4 ^d	49.8 \pm 1.8 ^b
4. <i>Cc.venom</i> _{i.p} + <i>Ao</i> extract _{i.p} group	90.8 \pm 5.1 ^b	58.0 \pm 4.5 ^d	33.7 \pm 1.4 ^b	67.2 \pm 1.7 ^a	75.7 \pm 3.5 ^a
5. <i>Ao</i> extract _{oral} then 2h + <i>Cc.venom</i> _{i.p} group	89.6 \pm 2.1 ^b	75.5 \pm 3.6 ^{ab}	32.5 \pm 1.2 ^b	59.0 \pm 1.8 ^{bc}	77.6 \pm 0.9 ^a
6. <i>Cc.venom</i> _{i.p} + <i>Antivenom</i> _{i.p} group	77.8 \pm 3.6 ^c	77.0 \pm 4.6 ^a	39.5 \pm 1.4 ^a	64.5 \pm 1.8 ^{ab}	42.2 \pm 1.6 ^c

Means within the same column in each category carrying different litters are significant at (P \leq 0.05).

Where

C.c venom _____ **Cerastes cerastes venom**
Ao extract _____ **Alkanna orientalis extract**
I.p _____ **intraperitoneal**

IV. DISCUSSION

Concerning to the effect on some kidney function parameter, serum calcium and serum cholinesterase. Our results showed that *Alkanna orientalis* extract did not cause any significant change in all previous determined parameters when given as a single intraperitoneal dose to albino rats as compared with control group. While, 2 hours after *Cerastes cerastes* venom injection lead to a significant increasing in serum creatinin, urea and uric acid also, a significant decrease in serum cholinesterase and calcium as compared with control group.

On the same pattern, anti-venom and *Alkanna orientalis* extract when given as antidote for *Cerastes cerastes* venom toxicity according to the program of treatment, succeed to ameliorate the toxicity of *Cerastes cerastes* venom on kidney function parameters.

The renal toxicity occurred after *Cerastes cerastes* venom injection agree with [35], as they documented acute renal failure with vascular lesions and tubular necrosis in renal cortex after different snake bites. In the present study, the elevation in serum uric acid, urea and creatinine levels indicate renal impairment. The increasing of urea and creatinine levels that observed in this study, reflect impairment of renal function [36] and [37]. Also, a similar observation was reported in rats after administration of various viper venoms were reported [38].

The decreasing in serum cholinesterase levels in rats following *Cerastes cerastes* venom injection is in agreement with [39] as they documented that Phospholipases A₂ is the major components of *Cerastes cerastes* venom and it exhibit neurotoxicity. Also, [40] reported that PLA₂ exhibit a presynaptic neurotoxicity.

According to the present study results, the significant decreasing in calcium level after *Cerastes cerastes* venom injection is in agreement with [41], as they illustrated the effect of sub lethal doses of *Cerastes cerastes* and *Macrovipera mauritanica* venom after 1, 3, 6 and 24 hours of envenomation, and the results showed that the venom caused a significant decreasing in calcium level in rabbit.

The protective action of *Alkanna orientalis* treatment is explicable in terms of their capacity for trapping free radicals and their stabilizing effect on the cytoplasmic membranes, they promotes protein synthesis, helps in regenerating kidney tissues[42]. As well as, *Alkanna orientalis* considered a powerfull anti-oxidant as it rich with Flavanoids that considered as one of the foremost plant components that work against PLA₂, lipooxygenase. The flavonoid was reported as strongly inhibits the group II PLA₂ from *Vipera* [21]. As well as , *Alkanna orientalis* rich source of rosmarinic acid, alpha - linolenic acid (ALA) and gamma- linolenic acid

(GLA) that acts as powerful anti-inflammatory and anti-oxidant against *Cerastes cerastes* venom renal toxicity [17] and [19].

The protective effect of *Alkanna orientalis* extract against the significant decreasing in cholinesterase and the cholinergic effect of *cerastes cerastes* venom is in full agreement with [43] as they described the protective effect of *Echium humile* extract (family: *Boraginacea*) against malathion neurotoxicity and against the significant decreasing in cholinesterase as a result of malathion cholinergic toxicity .

As well as, *Alkanna orientalis* rich with flavonoids that considered as one of the most plant ingredients that work against PLA2, as flavonoid is strongly inhibits the group II PLA2 from *Vipera* that responsible for the major *Cerastes cerastes* venom toxicity so, it able to antagonize the hazardous neurotoxic effect of the crude venom as well as *Cerastes cerastes* effect on serum calcium level [21].

Concerning to the effect on glucose and lipogram parameters (Total cholesterol, triglyceride, HDL-cholesterol and LDL-cholesterol) the obtained results revealed that, *Alkanna orientalis* extract when given as a single intraperitoneal dose did not cause any significant change in all determined lipogram parameters and glucose with the exception of triglyceride showed a significant decreasing as compared with control group. While, *Cerastes cerastes* venom according to the same program of treatment caused a significant decreasing in all lipogram parameters, Total cholesterol, triglyceride, HDL-cholesterol and LDL-cholesterol as well as, glucose level when compared with control group.

On the other hand , *Alkanna orientalis* extract and antivenom when given as antidote to *Cerastes cerastes* venom according to the program of treatment, succeed to modulate the *Cerastes cerastes* venom toxic effect and returned the lipogram and glucose parameter to about the normal value.

The significant hypoglycemic effect that observed after 2 hours from injection of the venom is in agreement with [44], [45] and [46], as the reduction of blood glucose level reflects a disturbance in carbohydrate metabolism, which could be referred to insulin-releasing effect of some venom components. As well as, the decreasing in serum cholesterol, triglycerides, HDL-cholesterol and LDL-cholesterol levels in rats following *Cerastes cerastes* venom injection, these findings are in agreement with the other investigators [47] and [45]. They reported that, a significant decreasing in serum cholesterol and triglyceride levels noted in laboratory animals that injected with snake venoms, suggested that the snake venom able to mobilize lipids from adipose tissues then splitting by lipolytic enzymes, that lead to free fatty acids liberation.

We suggested that *Cerastes cerastes* venom toxicity responsible for production of oxidative stress, which is the key contributor in hepatic injury and it known to produce reactive oxygen species (ROS) that is responsible for significant change in lipid profile and hepatic dysfunction.

The ameliorative effect of *Alkanna orientalis* that observed in this study against the hypoglycemic and hypolipemic effect of *Cerastes cerastes* venom explained by that, *Alkanna orientalis* rich with alpha-linolenic acid (ALA), gamma- linolenic acid (GLA) and rosmarinic acid that acts as antioxidant and anti inflammatory against *Cerastes cerastes* venom toxicity [17] and [19] .

ALA and GLA, act as anti-inflammatory and lipid lowering potential, it also cause vasodilation. In addition, GLA are important constituents of membrane phospholipids, including the mitochondrial membrane, where they enhance the integrity and the fluidity of the membrane [48]. As well as, dietary supplementation with GLA alone has yielded variable results on circulating lipid levels [49] and [50].

As well as, *Alkanna orientalis* considered a powerfull antioxidant as it rich with Flavanoids that considered as one of the foremost plant components that work against PLA2, lipooxygenase. They possess anti-inflammatory, anti-hypertensive, hypocholesterolemic, antiallergic, and enzyme inhibiting properties. The flavonoid strongly inhibits the group II PLA2 from *Vipera* that responsible for the major toxicity of *Cerastes cerastes* venom so, it able to antagonize the hazardous toxic effect of the crude venom [21].

V –CONCLUION

The natural medical herbs are showing a new area for development of better therapeutic and prophylaxis agents against expected envenomation by *Cerastes cerastes* venom toxicity. The alcoholic extract of *Alkanna orientalis* has many benefits, as it is cheap, stable at room temperature and able to antagonize the hazardous toxic effect of *Cerastes cerastes* venom. Therefore, we recommend the usage of *Alkanna orientalis* extract as initial assist in *Cerastes cerastes* bitted victims therapy to minimize mortality and morbidity. As well as, this extract can be used as a prophylaxis for researchers who work in the fields where *Cerastes cerastes* are abundant.

REFERENCES

- [1] **Chippaux JP, Lang J, Amadi-Eddine S, Fagot P, Le Mener V. (1999).** Short report: treatment of snake envenomations by a new polyvalent antivenom composed for highly purified F(ab')₂. Results of a clinical trial in northern Cameroon. *Am J Trop Med Hyg* –1018 - 1017:(6) 61, 1999.
- [2] **Theakston RD, Warrell DA, Griffiths E.(2003).** Report of a WHO workshop on the standardization and control of antivenoms. *Toxicon* 2003, 41(5):541–557.
- [3] **Frobert, Y., Créminon, C., Cousin, X., Rémy, M.H., Chatel, J.M., Bon, S., Bon, C., Grassi, J. (1997).** Acetylcholinesterases from Elapidae snake venoms: biochemical, immunological and enzymatic characterization. *Biochim Biophys Acta*. 1339: 253-67.
- [4] **Syed Moied Ahmed, Mohib Ahmed, Abu Nadeem, Jyotsna Mahajan, Adarash Choudhary, and Jyotishka Pal. (2008).** Emergency treatment of a snake bite: *Pearls from literature J Emerg Trauma Shock*. 2008 Jul-Dec; 1(2): 97–105.
- [5] **Anai K, Sugiki M, Yoshida E, Maruyama M. (2002).** Neutralization of a snake venom hemorrhagic metalloproteinase prevents coagulopathy after subcutaneous injection of *Bothrops jararaca* venom in rats. *Toxicon* 2002; 40: 63–68.
- [6] **Kebir-Chelghoum H, Laraba-Djebari F. (2017).** Cytotoxicity of Cerastes cerastes snake venom: Involvement of imbalanced redox status. *Acta Trop* 2017;173:116-24.
- [7] **Ibrahim AM, Elsefi TT, Ghanem M, Fayed AM, Shaban NA(2017).** A horned viper bite victim with PRES. *Case reports in neurological medicine*. 2017; Article ID 1835796, 6 pages
- [8] **Leon G, Valverde JM, Rojas G, Lomonte B, Gutierrez JM.(2000).** Comparative study on the ability of IgG and Fab sheep antivenoms to neutralize local hemorrhage, edema and myonecrosis induced by *Bothrops asper* (terciopelo) snake venom. *Toxicon* 2000;38:233–244.
- [9] **Girish KS, Kemparaju K.(2006).** Inhibition of *Naja naja* venom hyaluronidase role in the management of poisonous bite. *Life Sci* 2006; 78: 1433–1440.
- [10] **Girish KS, Kemparaju K.(2007)** The magic glue hyaluronan and its eraser hyaluronidase: a biological overview. *Life Sci*,C 2007; 80(21): 1921–1943.
- [11] **De smet P A (1997):** The role of plant derived drugs and herbal medicines in health care. *Drugs* ;54:801-840 .
- [12] **Hasani-Ranjbar S, Larijani B, Abdollahi M.(2009).** A systemic review of potential herbal sources of future drugs effective in oxidant related diseases. *Allergy Drug targets* 2009; 8(1): 2 -10.
- [13] **Aung HT, Nikai T, Niwa M, Takaya Y.(2010).** Rosmarinic acid in inhibits *Argusia argentea* snake venom-induced hemorrhage. *J Nat Med* 2010; 64(4): 482 – 486.
- [14] **Sekhar J, Penchala G, Sudarsanam G, Prasad GP.(2011).** Ethnic information on treatment for snake bites in kadapa district of Andhra Pradesh. *Life Science Leaflet* 2011; 12; 368 – 375.
- [15] **Gomes A, Das R, Sarkhel S, Mishra R, Mukherjee S, Bhattacharya S.(2010).** Herbs and herbal constituents active against snakebite. *Indian J Exp Biol* 2010; 48(9):865-78.
- [16] **Abbaszadeh S, Radjabian T, Taghizadeh M, Fazeli F, Salmaki Y. (2011).** Characterization of fatty acids in different organs of some Iranian *Echium* plants. *J Med Plants Res* 2011;5(19):4814-21.
- [17] **Katrin K, Degen C, Jaudszus A, and Jahreis G.(2012).** Searching for health beneficial n-3 and n-6 fatty acids in plant seeds. *Eur J Lipid Sci Technol*. 2012 Feb; 114(2): 153–160.
- [18] **Petersen M, Simmonds MS.** Rosmarinic acid. *Phytochem* 2003; 62(2): 121– 125.
- [19] **Ticli FK, Hage LI, Cambraia RS, Pereira PS, Magro AJ, Fontes MR, Stábeli RG, Giglio JR, França SC, Soares AM, Sampaio SV.(2005).** Rosmarinic acid, a new snake venom phospholipase A2 inhibitor from *Cordia verbenacea* (Boraginaceae): antiserum action potentiation and molecular interaction. *Toxicon* 2005; 46(3):318-27.
- [20] **El Sohly HN, El-Ferally FS, Joshi AS, Walker LA.(1997).** Antiviral flavonoids from *Alkanna orientalis*. *Planta Medica* 1997; 63: 384.
- [21] **Gopi K, Renu K, Jayaraman G.(2014).** Inhibition of *Naja naja* venom enzymes by the methanolic extract of *Leucas aspera* and its chemical profile by GC-MS. *Toxicol Rep* 2014;1:667-673.
- [22] **Ramakrishnan M A .** Determination of 50% endpoint titer using a simple formula. *World J Virol*: 2016; 5: 85–86.
- [23] **Sukhdev SH, Suman PS, Gennaro L, Dev DR. (2008).** Extraction technologies for medicinal and aromatic plants. *United Nation Industrial Development organization and the international center for Science and High Technology*. 2008;116.
- [24] **Henry, R.G. (1974):** Clinical Chemistry. Chemicals and Technique. (2nd Ed), Harper, New york . P.257.

- [25] **Patton, C. and Crouch, G. (1977)**: Enzymatic colorimetric determination of urea. *Anal .Chem .*;49 :464 - 469.
- [26] **Trinder P (1969b)**: Enzymatic method for uric acid determination. *Ann. Clin. Biochem.*, 3(6):29-30.
- [27] **Tietz NW. (1994)**. Fundamentals of Clinical Chemistry: 2 nd ed. *Tietz NW, editor, 1994, pp 692.*
- [28] **Szasz , G (1968).** effect of drugs on clinical lab . clinical chem . *Acta 19 . 191*
- [29] **Trinder, P. (1969a)**: Enzymatic method for glucose determination. *Ann. Clin. Biochem*; 3 (6):29-30.
- [30] **Richmond, W. (1973)**: Preparation and properties of cholesterol oxidase. *Amer. J. Clin. Chem*; 19 :1350-1356.
- [31] **Fossati, P.; and Prencipe, L. (1990)**: Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide. *Clin. Chem*; 6 (28):2077-2079.
- [32] **Stein, E.A. (1986)**: Textbook of Clinical Chemistry. *Nwtietz, ed., W.B Saunders, Philadelphia. PP. 879-886.*
- [33] **Duncan DB.** Multiple range and multiple F tests. *Biometrics 1955; 11(1): 1-42.*
- [34] **Armitage P, Berry G.(1987)**. Statistical Methods in Medical Research. *Boston MA: Blackwell Scientific, Oxford, UK, 1987; 559.*
- [35] **Tilbury RC, Madkour MM, Saltissi D., Suleiman M. (1987)**. Acute renal failure following the bite of Burton's Carpet Viper *Echis coloratus* Gunther in Saudi Arabia: *Case report review. Saudi. Med. J. 8: 87-95.*
- [36] **Mora, L.O; Antunes, L.M.G; Francescato, H.D.C and Bianchi, M.L.P.(2003)**: The effects of oral glutamine on cisplatin-induced nephrotoxicity in rats. *Pharmacol. Res.*, 47: 517-522).
- [37] **Yousef, M.I; Awad, T.I and Mohamed, E.H (2006)**: Deltamethrininduced oxidative damage and biochemical alterations in rat and its attenuation by vitamin E. *Toxicology, 111: 240-247, (2006).*
- [38] **Schneemann M, Cathomas R, Laidlaw ST, El Nahas AM, Theakston RD, Warrell DA (2004)**. Lifethreatening envenoming by the Saharan horned viper(*Cerastes cerastes*) causing microangiopathic haemolysis, coagulopathy and acute renal failure. *Clinical cases and review. 97(11): 717-727.*
- [39] **Soares AM, Giglio JR.(2003)**. Chemical modifications of phospholipases A2 from snake venoms: *effects on catalytic and pharmacological properties. Toxicon 2003;42(8), 855-868.*
- [40] **Kini RM. (2003)**. Excitement ahead: structure, function and mechanism of snake venom phospholipase A2 enzymes. *Toxicon 2003;42(8): 827–840.*
- [41] **Fahmi L, B. Makran A, L. Boussadda B, M. Lkhider A, N. Ghalim C (2014)**. Study of biochemical and hematological changes after envenomation by viper *Cerastes cerastes* and *Macrovipera mauritanica* in rabbit. *Abstracts from 21st Meeting of the French Society of Toxinology (SFET) / Toxicon 91 (2014) 164e184.*
- [42] **Naglaa ,M .El-Lakkany; Olfat .A. Hammam; Walaa ,H. El- Maadawy; Afkar A ,Badawy; Afaf ,A, Ain-Shoka and Fatma, A. Ebeid(2012)**: Anti-inflammatory/anti-fibrotic effects of the hepatoprotective silymarin and the *Schistosomicide praziquantel* against *Schistosoma mansoni*-induced liver fibrosis . *Parasites & Vectors, 5:9*
- [43] **Diab AA, Dahmy SI , Soliman. SAI and Nasser M M.(2013)**. The protective role of *Echium humile* extract against toxicity induced by insecticide (malathion) in male albino rats. *Journal of American Science 2013; 9 (11):259-267.*
- [44] **Abu-Sinna G, AL-Zahaby AS, Abd EL-Aal A, Abd- EL-Baset, Soliman NA (1993)**. The Effect of the viper *Cerastes cerastes cerastes* venom and venom fractions on carbohydrate metabolism. *Toxicon, 31(6): 791- 801.*
- [45] **Al-Jammaz I (2002)**. Effects of envenomation by *Creastes vipera* crude venom plasma and tissue metabolites of rats. *Kawait, J. Eng. 29(1): 111-119.*
- [46] **Al-Sadoon MK, Abdel Moneim AE, Diab MS, and Amira A, Bauomy AA(2013)**. Hepatic and renal tissue damages induced by *Cerastes cerastes gasperetti* crude venom. *Life Science J. 10(4):191-197.*
- [47] **Abdel-Nabi IM, Raafat A, El-Shamy HI (1997)**. Biological effects of intraperitoneal injection of rats with the venom of the snake *Echis carinatus*. *Egypt. J. Zool. 29: 195-205.*
- [48] **Horrobin DF.(1992)**. The use of gamma-linolenic acid in diabetic neuropathy. *Agents Actions Suppl 1992;37:120-144.*
- [49] **Graham J, Franks S, Bonney RC.(1994)**. In vivo and in vitro effects of gamma-linolenic acid and eicosapentaenoic acid on prostaglandin production and arachidonic acid uptake by human endometrium. *Prostaglandins Leukot Essent Fatty Acids 1994;50:321-329.*
- [50] **von Schacky C.(2000)**. n-3 fatty acids and the prevention of coronary atherosclerosis. *Am J Clin Nutr 2000;71:224S-227S.*